



Modeling the effect of surface washing treatment on inactivation of spoilage bacteria and shelf life extension of fresh fish



Theofania Tsironi, Ioanna Semenoglou, Athina Ntzimani, Efimia Dermesonlouoglou, Petros Taoukis

School of Chemical Engineering, National Technical University of Athens, Greece (taoukis@chemeng.ntua.gr)

Introduction

The short shelf life and perishability of fish products is a commercial drawback and methods of extension are being investigated. New minimal and nonthermal food processing methods are sought by the industry in the pursuit of producing better quality fish products with extended shelf life with retention of nutritional and sensory properties (Tsironi et al., 2014 and 2019; Tsironi and Taoukis, 2019). Several studies have been conducted recently on the efficacy of washing and sanitizing treatments in reducing microbial populations on food products. Limited work on the effect on fish has been published and no industrial scaling-up has been reported (Thi et al., 2015).

The objective of the study was to investigate and mathematically model the effect of surface decontamination of fresh fish using alternative organic acids on the quality and shelf life during refrigerated storage.

Materials & Methods

Marine cultured gilthead seabream (*Sparus aurata*) were stored isothermally at 0°C for 6 days after harvesting. Fish was gutted manually and immersed in water for 0-10 min. The incorporation of natural organic acids (lactic acid, citric acid) at different concentrations (0-200 ppm) for times 0-10 min during gutting was tested for its efficacy to reduce initial microbial load and prolong shelf life.

Control (treated with water) and organic acid treated samples were afterwards stored under controlled isothermal conditions (0-10°C) for shelf life testing. Quality assessment was based on microbiological analysis (total viable count, *Pseudomonas* spp., *Enterobacteriaceae* spp., lactic acid bacteria, H₂S-producing bacteria, etc), pH, colour, texture measurement and sensory scoring. A sensory score of 5 was taken as the average score for minimum acceptability.

Results

Initial surface decontamination (up to 2 logcfu/g for total viable count, *Pseudomonas* spp., *Enterobacteriaceae* spp. and H₂S-producing bacteria, depending on species and processing conditions) by the addition of organic acids in the washing water was observed (Figures 1 a-c).




Figure 1. Results after washing with lactic acid or citric acid at concentrations 0-200 ppm for 0-10 min on (a) Total viable count, (b) *Pseudomonas* spp. and (c) *Enterobacteriaceae*, in whole gilthead seabream.

Acknowledgment

The research was funded by the Operational Programme for Fisheries and Maritime 2014-2020-Greece, Priority 2, Measure 3.2.1, Article 47. (MIS 5010939-Development and application of novel methods for fish harvesting and processing for quality improvement and shelf life extension)

Microbial load reduction was increased for higher washing solution concentrations and longer treatment. Higher reduction of the initial microbial load was observed after treatment with citric acid for TVC, *Pseudomonas* spp. and H₂S-producing bacteria and with lactic acid solution for *Enterobacteriaceae* spp.

Mathematical models were developed for the inactivation of spoilage bacteria as a function of treatment conditions and the concentration of acid in the washing water. The experimental data were adequately described by Equation (1):

$$\log\left(\frac{N}{N_0}\right) = a * \sqrt{C - b} * e^{-d*t} \quad (1)$$

where N_0 and N are the initial and final (after treatment) microbial load, C is % (w/v) washing solution concentration, t is treatment (min) and a , b , d are constants.

Microbial growth during subsequent refrigerated storage of untreated (Control) and treated fish was modeled using the Baranyi Growth Model. Limit of sensory shelf life of gutted fish (score 5 by the sensory panel for overall impression) coincided with a *Pseudomonas* spp. level of 10⁷ cfu/g at all tested storage temperatures (0-10°C). Based on the values at the end of the shelf life of the studied indices and the temperature dependence of their rate constants expressed by the Arrhenius kinetics, simple equations for shelf life calculation can be used (Equation 2).

$$t_{SL} = \frac{\log N_L - \log N_0}{k_{ref} \cdot \exp\left[\frac{-E_a}{R} \cdot \left(\frac{1}{T} - \frac{1}{T_{ref}}\right)\right]} \quad (2)$$

where t_{SL} is the shelf life (d) of gilthead seabream, $\log N_L$ is the limit *Pseudomonas* spp. load (7 log cfu/g), $\log N_0$ is the initial *Pseudomonas* spp. load, k_{ref} is the rate constant of *Pseudomonas* spp. growth at a reference temperature T_{ref} (4°C), E_a is the activation energy of *Pseudomonas* spp. growth, R is the universal gas constant.

Based on these calculations, the shelf life of gutted gilthead seabream for different treatment conditions is presented in Table 1.

Table 1. Shelf life (days) of gutted gilthead seabream for different processing and storage conditions.

Processing / storage conditions	Control	Citric acid (100ppm/5 min)	Citric acid (200 ppm/10 min)
0°C	12	13	16
5°C	6	7	8
10°C	4	5	6

Conclusions

The results of the study indicated that the application of washing treatment led to improved quality stability during subsequent refrigerated storage and shelf life extension. Initial surface decontamination up to 2.0 logcfu/g by the addition of organic acids in the washing water may result in 2-4 days shelf life extension of gutted fish at 0°C storage. Shelf life extension of fish could open new distant markets currently inaccessible to fresh fish products and allow the use of higher temperatures (e.g. 5°C) in the cold chain of seafood which would significantly reduce energy and food waste.

References

- Thi, A.N.T., Sampers, I., Haute, S., Nguyenc, B.L., Heyndrickx, M., Devlieghere, F. 2015. Decontamination of Pangasius fish (*Pangasius hypophthalmus*) with chlorine or peracetic acid in the laboratory and in a Vietnamese processing company. *Int J Food Microbiology*, 208, 93-101.
- Tsironi T., Anjos L., Pinto P.I.S., Dimopoulos G., Santos S., Santa C., Manadas B., Canario A., Taoukis P., Power D. (2019). High pressure processing of European sea bass (*Dicentrarchus labrax*) fillets and tools for flesh quality and shelf life monitoring. *J Food Engi*, 262, 83-91.
- Tsironi T., Maltezou I., Tsevdou M., Katsaros G., Taoukis P.S. (2015). High pressure cold pasteurization of gilthead seabream fillets: Selection of process conditions and validation of shelf-life extension. *Food and Bioprocess Technology: An International Journal*, 8, 681-690.
- Tsironi T., Taoukis P. (2019). Advances in conventional and nonthermal processing of fish for quality improvement and shelf life extension. *Reference Module in Food Science*. Elsevier, pp. 1-7.

